
Variable Metric Stochastic Approximation Theory

Peter Sunehag*
Peter.Sunehag@nicta.com.au
NICTA, Locked Bag 8001
Canberra ACT 2601
Australia

Jochen Trumpf
Jochen.Trumpf@anu.edu.au
Dept. of Information Engr.
Australian National Univ.
Canberra, Australia

S.V.N. Vishwanathan
vishy@stat.purdue.edu
Statistics Department
Purdue University
West Lafayette, IN

Nicol N. Schraudolph
schraudolph@adaptivetools.com
adaptive tools AG
Canberra ACT 2602
Australia

Abstract

We provide a variable metric stochastic approximation theory. In doing so, we provide a convergence theory for a large class of online variable metric methods including the recently introduced online versions of the BFGS algorithm and its limited-memory LBFGS variant. We also discuss the implications of our results for learning from expert advice.

1 Introduction

We begin by introducing online optimization and in particular stochastic gradient descent methods.

1.1 Online Optimization

There exists an abundance of applications that can lead us to online optimization problems where we are trying to find the minimum of a data-dependent function $C : \mathbb{R}^n \rightarrow \mathbb{R}, w \mapsto C(w) = C(w, Z)$, where Z represents data or a probability distribution that generates data. During an online optimization procedure, a sequence of parameters $w_t, t = 1, 2, \dots$ is created by using an update rule for how to define w_{t+1} given the earlier parameters and the new data, z_t , that has arrived at time t . Given a sequence of data samples z_1, \dots, z_t drawn from a fixed distribution, we will use the notation $C_t(\cdot) = \frac{1}{t} \sum_{i=1}^t C(\cdot, z_i)$ for the empirical objective which is the average of online/stochastic/instantaneous objectives $C(\cdot, z)$. We will refer to $C(\cdot) = \mathbb{E}_z C(\cdot, z)$ as the true objective.

*Also an adjunct at the second author's affiliation.

Appearing in Proceedings of the 12th International Conference on Artificial Intelligence and Statistics (AISTATS) 2009, Clearwater Beach, Florida, USA. Volume 5 of JMLR: W&CP 5. Copyright 2009 by the authors.

1.2 Stochastic Gradient Descent

If one has defined a metric κ on the parameter space by supplying a dot product, and if the online objectives are differentiable, then we can use the gradients ∇^κ with respect to that metric to define the update equation

$$w_{t+1} = w_t - a_t \nabla_w^\kappa C(w_t, z_t), a_t > 0. \quad (1)$$

If one uses metrics defined by different dot products for different t , then one can let ∇ denote the gradient with respect to the standard Euclidean dot product and instead let the updates take the form

$$w_{t+1} = w_t - a_t B_t \nabla_w C(w_t, z_t), \quad (2)$$

where the B_t are positive definite and symmetric matrices. One example of considering variable metrics is the study of information geometry (Amari and Nagaoka, 1993), where the Fisher information matrix is used to define a metric tensor on a family of probability distributions. More specific examples will be provided in Section 2 below.

1.3 Outline and Summary

We investigate the theoretical foundations for using online updates that include scaling matrices, that is stochastic gradient descent where the gradients are taken with respect to time-varying metrics. Among other results, this provides a convergence proof for a large class of variable metric methods including the recent online (L)BFGS algorithm (Schraudolph et al., 2007). In Section 3 we employ the Robbins-Siegmund theorem to prove $O(1/t)$ convergence in function values $C(w_t)$ for a class of functions that is useful in machine learning. This is the best possible rate (Bottou and LeCun, 2005; Amari, 1998; Murata, 1998), limited only by the rate at which information arrives. Under weaker assumptions we show almost sure convergence without rate for a larger class. Our results extend

those in Bottou and LeCun (2005) by not demanding that the metrics converge to an asymptotic metric.

We first introduce motivating application areas in Section 2. Then in Section 3 we both provide a background in stochastic approximation theory and our new theorems. We apply our theorems to prove convergence for online BFGS in Section 4. Furthermore, we consider implications in the area of learning from expert advice in Section 5. The paper concludes with a discussion in Section 6.

2 Examples

In this section we provide different objective functions and describe the problem settings they appear in.

2.1 Online Risk Minimization

The goal of many machine learning algorithms is to minimize the risk $\mathbb{E}_z l(\langle w, x \rangle, y)$, where the expectation is taken with respect to a fixed but unknown probability distribution Z that generates instance-label pairs $z := (x, y)$. The loss l is a non-negative convex function of the parameters w and measures the discrepancy between the labels y and the predictions arising from x and w via their inner product $\langle w, x \rangle$ (often the Euclidean dot product if $x, w \in \mathbb{R}^k$).

In the absence of complete knowledge about the underlying distribution, an empirical sample $Z = \{(x_i, y_i), i = 1, \dots, n\}^1$ is often used to minimize the regularized empirical risk

$$\frac{c}{2} \|w\|^2 + \frac{1}{n} \sum_{i=1}^n l(\langle w, x_i \rangle, y_i) \quad (3)$$

where the L_2 -regularization term $\frac{c}{2} \|w\|^2$ is introduced for well-posedness. With larger n one can use smaller $c > 0$. The true regularized risk that is estimated by (3) is $\frac{c}{2} \|w\|^2 + \mathbb{E}_z l(\langle w, x \rangle, y)$.

Batch optimization algorithms, including quasi-Newton and bundle-based methods, are available and widely used to minimize (3), but they are computationally expensive. Gradient-based batch methods may also fail to converge if the loss is non-smooth. Therefore, online optimization methods that work with small subsamples of training data have received considerable attention recently (Kivinen and Warmuth, 1997; Schraudolph, 2002; Azoury and Warmuth, 2001; Shalev-Shwartz et al., 2007; Schraudolph et al., 2007).

In the online setting, we replace the objective (3) with approximations based on subsets (“mini-batches”) of

¹With some abuse of notation we use Z to represent either a data set or a distribution that generates data.

the samples:

$$\frac{c}{2} \|w\|^2 + \frac{1}{b} \sum_{(x,y) \in Z_t} l(\langle w, x \rangle, y), \quad (4)$$

where $Z_t \subset Z$ with $|Z_t| = b \ll n$. Furthermore, in online learning we can consider using $c = 0$ and thereby aiming directly at minimizing the true objective $\mathbb{E}_z l(\langle w, x \rangle, y)$. During online optimization, a sequence of parameters w_t , $t = 1, 2, \dots$ arises from an update rule that computes w_{t+1} from the previous state and the new information at time t . In addition to alleviating the high computational cost of batch methods, the online setting also arises when the data itself is streaming, that is, we are receiving partial information about $C(\cdot)$ in a sequence of small packages.

Including second-order information in the online optimization procedure can accelerate the convergence (Schraudolph et al., 2007). This is particularly true in a setting where only one pass through a dataset is performed. Bottou and LeCun (2005) and Murata (1998) point out that minimizing the empirical objective is different from minimizing the true objective, and show that the result of an online second-order gradient descent procedure can be as close to the true optimum as the minimum of the empirical objective.

2.2 Filtering

The goal in filtering is to separate the signal from the noise in a stream of data. Kalman algorithms in particular use the Euclidean distance (sum-squared loss) and track the minimizer of the empirical objectives $C_t(w) = \sum_{i=1}^t (y_t - w \cdot x_t)^2$. The inverse of the Hessian of C_t is used as B_t , $a_t = 1$ and $w_0 = 0$ for the update in (2). B_{t+1} is found from B_t with an update whose cost is order n^2 . The result is that $w_t = \operatorname{argmin}_w C_t(w)$. Therefore, if we have a fixed distribution the sequence will converge to the optimal parameters.

The same algorithm can be extended to a more general setting where the sum-squared loss is replaced by arbitrary convex functions. The resulting algorithm was called the online Newton-step algorithm by Hazan et al. (2007) and described as an approximate “follow the leader” algorithm, *i.e.*, an algorithm that approximately follows the optimum that the Kalman filter tracks exactly for the sum-squared loss.

2.3 Learning from Expert Advice with Bregman Divergences

The general convex optimization framework by Zinkevich (2003) that Hazan et al. (2007) worked in is also related to the expert advice framework by Azoury and Warmuth (2001). In the expert advice framework one encounters a sequence of loss functions L_t

($L_t(w) = C(w, z_t)$ here) and one wants to perform the implicitly defined update

$$w_{t+1} = \underset{w}{\operatorname{argmin}} \Delta_H(w, w_t) + a_t L_t(w) \quad (5)$$

where Δ_H is the Bregman divergence

$$\Delta_H(p, q) = H(p) - H(q) + \nabla H(q) \cdot (p - q) \quad (6)$$

defined by the differentiable convex function H . The squared Euclidean distance is an example of a Bregman divergence, corresponding to $H(p) = \frac{1}{2} \|p\|_2^2$. The goal for Azoury and Warmuth (2001), as it was for Zinkevich (2003), was to have a small total accumulated loss $\sum_{j=1}^t L_j(w_j)$ and in particular small regret, $\sum_{j=1}^t L_j(w_j) - \min_w \sum_{j=1}^t L_j(w)$. To derive an explicit update, Azoury and Warmuth (2001) differentiate the expression to be minimized

$$\begin{aligned} \nabla(\Delta_H(w, w_t) + a_t L_t(w)) = \\ h(w) - h(w_t) + a_t \nabla L_t(w) \end{aligned} \quad (7)$$

and by using the approximation $\nabla L_t(w) \approx \nabla L_t(w_t)$, they arrive at the updates

$$w_{t+1} = h^{-1}(h(w_t) - a_t \nabla L_t(w_t)). \quad (8)$$

This reduces to online gradient descent, $w_{t+1} = w_t - \nabla L_t(w_t)$, for the case $h(w) = w$, *i.e.*, when $H(w) = \frac{1}{2} \|w\|_2^2$.

Azoury and Warmuth (2001) in particular consider loss functions of a special form, namely the case where $L_t(w) = \Delta_G(x_t \cdot w, g^{-1}(y_t))$ where $g = \nabla G$ is an increasing continuous function from \mathbb{R} to itself and where $(x_t, y_t) = z_t$ is an example. This results in simpler updates since it implies that $\nabla L_t(w) = (g(w \cdot x_t) - y_t)x_t$. Note that, given a parametrization of \mathbb{R}^d , a one-dimensional transfer function g can be used to define a d -dimensional continuous bijection by applying it coordinate-wise.

It is interesting to compare (8) to a reparametrization where one makes a coordinate change $\gamma = h(w)$. Then (8) can be written as

$$h(w_{t+1}) = h(w_t) - a_t \nabla_w L_t(w_t). \quad (9)$$

The one obstacle to identifying (9) with

$$\gamma_{t+1} = \gamma_t - a_t \nabla_\gamma \tilde{L}_t(\gamma_t) \quad (10)$$

where $\tilde{L}_t(\gamma) = L(h^{-1}(\gamma))$, is that the gradient of L_t is taken with respect to w . We know that $\nabla_\gamma \tilde{L}_t(\gamma_t) = B_t \nabla_w L_t(w_t)$, where B_t is the inverse Hessian of H at w_t , in other words the inverse Jacobian of h .

A reason for doing a reparametrization is that the loss functions L_t might not satisfy the conditions that are

needed for convergence of stochastic gradient descent, but it can be possible to find a transfer function h such that \tilde{L}_t does. In that sense, the update (8) can be a way of trying to do this even if we only have the gradients $\nabla_w L_t$ and do not want to calculate the matrices B_t . Our main theorems will tell us when it is acceptable to omit B_t , and thereby use an SGD update with respect to varying metrics.

3 Stochastic Approximation Theory

Robbins and Monro (1951) proved a theorem that implies convergence for one-dimensional stochastic gradient descent; Blum (1954) generalized it to the multivariate case. Robbins and Siegmund (1971) achieved a stronger result of wider applicability in supermartingale theory. Here we extend the known convergence results (Bottou and LeCun, 2005) in two ways: a) We prove that updates that include scaling matrices with eigenvalues bounded by positive constants from above and below will converge almost surely; b) under slightly stronger assumptions we obtain a $O(1/t)$ rate of convergence in the function values.

3.1 The Multivariate Robbins-Monro Procedure

Suppose that $\nabla C = f : \mathbb{R}^k \rightarrow \mathbb{R}^k$ is an unknown continuous function that we want to find a root of. Furthermore, suppose that there is a unique root and denote it by w^* . Given an initial estimate w_1 , the procedure constructs a sequence of estimates w_t such that $w_t \rightarrow w^*$ as $t \rightarrow \infty$. For any random vector X let $\mathbb{E}_t(X)$ be the conditional expectation given w_1, \dots, w_t . Given w_1, \dots, w_t , we assume that we observe an unbiased estimate Y_t of $f(w_t) = \nabla C(w_t)$, *i.e.*, $\mathbb{E} Y_t = f(w_t)$. Given Y_t and w_t we define w_{t+1} by

$$w_{t+1} = w_t - a_t Y_t, \quad (11)$$

where $a_t > 0$ for all t , $\sum a_t = \infty$ and $\sum a_t^2 < \infty$. The Y_t are assumed to be drawn from a family $Y(x)$ of random vectors defined for all $x \in \mathbb{R}^k$, and Y_t is distributed as $Y(w_t)$. To ensure that w_t converges to w^* almost surely it is sufficient to assume that there are finite constants A and B such that $\mathbb{E} \|Y(x)\|^2 \leq A + B \|x - w^*\|^2$ for all x , and that for all $\varepsilon > 0$

$$\inf\{(x - w^*)^T f(x) : \varepsilon < \|x - w^*\| < \varepsilon^{-1}\} > 0. \quad (12)$$

For instance, strictly convex functions satisfy (12). This classical convergence result is implied by the following theorem on almost positive supermartingales, which we will also use to prove our results:

Theorem 3.1 (Robbins and Siegmund, 1971)

Let (Ω, \mathcal{F}, P) be a probability space and $\mathcal{F}_1 \subseteq \mathcal{F}_2 \subseteq \dots$

a sequence of sub- σ -fields of \mathcal{F} . Let U_t, β_t, ξ_t and ζ_t , $t = 1, 2, \dots$ be non-negative \mathcal{F}_t -measurable random variables such that

$$\mathbb{E}(U_{t+1} \mid \mathcal{F}_t) \leq (1 + \beta_t)U_t + \xi_t - \zeta_t, \quad t = 1, 2, \dots \quad (13)$$

Then on the set $\{\sum_t \beta_t < \infty, \sum_t \xi_t < \infty\}$, U_t converges almost surely to a random variable, and $\sum_t \zeta_t < \infty$ almost surely.

3.2 Convergence for Updates with Scaling Matrices

Bottou and LeCun (2005) previously presented results on the rate of convergence in parameter space (using the Euclidean norm) of online updates with *convergent* scaling matrices, and remark that bounds on the eigenvalues of the scaling matrix will be essential to extending convergence guarantees beyond that. Since recent online quasi-Newton methods (Schraudolph et al., 2007) do not provide convergence of their scaling matrices, we now employ the Robbins-Siegmund theorem to prove that such updates are still guaranteed almost sure convergence, provided the spectrum of their (possibly non-convergent) scaling matrices is uniformly bounded from above by a finite constant and from below by a strictly positive constant.

Theorem 3.2 *Let $C : \mathbb{R}^n \rightarrow \mathbb{R}$ be a twice differentiable cost function with unique minimum w^* , and let*

$$w_{t+1} = w_t - a_t B_t Y_t, \quad (14)$$

where B_t is symmetric and only depends on information available at time t .

Then w_t converges to w^* almost surely if the following conditions hold:

- C.1** $(\forall t) \mathbb{E}_t Y_t = \nabla_w C(w_t)$;
- C.2** $(\exists K) (\forall w) \|\nabla_w^2 C(w)\| \leq 2K$;
- C.3** $(\forall \delta > 0) \inf_{C(w) - C(w^*) > \delta} \|\nabla_w C(w)\| > 0$;
- C.4** $(\exists A, B) (\forall t) \mathbb{E} \|Y_t\|^2 \leq A + BC(w_t)$;
- C.5** $(\exists m, M : 0 < m < M < \infty) (\forall t) mI \prec B_t \prec MI$, where I is the identity matrix;
- C.6** $\sum_t a_t^2 < \infty$ and $\sum_t a_t = \infty$.

Proof Since C is twice differentiable and has bounded Hessian (C.2) we can use Taylor expansion and the upper eigenvalue bound (C.5) to prove that

$$C(w_{t+1}) = C(w_t - a_t B_t Y_t) \leq \quad (15)$$

$$C(w_t) - a_t [\nabla_w C(w_t)]^T B_t Y_t + KM^2 a_t^2 \|Y_t\|^2$$

which implies, using (C.1) and (C.4), that

$$\mathbb{E}_t C(w_{t+1}) \leq C(w_t) + KM^2 a_t^2 [A + BC(w_t)] - \quad (16)$$

$$a_t [\nabla_w C(w_t)]^T B_t \nabla_w C(w_t).$$

If we let $U_t = C(w_t)$ and merge the terms containing U_t it follows that $\mathbb{E}_t U_{t+1} \leq$

$$U_t(1 + a_t^2 BKM^2) + AKM^2 a_t^2 - ma_t \|\nabla_w C(w_t)\|^2. \quad (17)$$

Since $\sum_t a_t^2 < \infty$ (C.6), the Robbins-Siegmund theorem can now be applied. We find that

$$\sum_t a_t \|\nabla_w C(w_t)\|^2 < \infty. \quad (18)$$

Since $\sum a_t = \infty$ (C.6) it follows from (C.3) that

$$\|(\nabla_w C(w_t))\|^2 \rightarrow 0 \quad (19)$$

and that $C(w_t) \rightarrow C(w^*)$ as $t \rightarrow \infty$. ■

Remark 3.3 *The assumption that C is twice differentiable is only needed for the Taylor expansion we use to obtain (15). If we have such a property from elsewhere we do not need twice-differentiability.*

3.3 Asymptotic Rates

Consider the situation described in Theorem 3.2. We now strengthen assumption (C.3), which demands that the function C is not so flat around the minimum that we may never approach it, to instead assuming that

$$\frac{C(w_t) - C(w^*)}{\|\nabla C(w_t)\|^2} \leq D < \infty \quad \forall t. \quad (20)$$

Condition (20) is implied by strong convexity if we know that the w_t tend to the optimum w^* . Since $\nabla C(w^*) = 0$ we can use first-order Taylor expansion of ∇C around w^* to approximate $\|\nabla C(w)\|^2$ by $(w - w^*)^T \nabla^2 C(w^*)(w - w^*)$.

We will also modify assumption (C.4) by setting $B = 0$. Theorem 3.2 guarantees (under the weaker conditions) that the procedure will almost surely generate a converging sequence which is therefore contained in some ball around w^* . This makes the new condition reasonable. Bottou (1998) contains a more elaborate discussion on what is there called "Global Confinement".

We need a result on what the expected improvement is, given the step size a_t , the uniform bound on the Hessian, and the uniform eigenvalue bounds. The key to achieving this is (15). This section's counterpart to (16) under the new conditions is

$$\mathbb{E}_t C(w_{t+1}) - C(w^*) \leq \quad (21)$$

$$[C(w_t) - C(w^*)](1 - a_t m D) + AKM^2 a_t^2.$$

We want to know the rate of the sequence $\mathbb{E}C(w_t) - \inf_{\tilde{w}} C(\tilde{w})$, and will use the fact that taking the unconditional expectation of (21) yields the result

$$\mathbb{E}C(w_{t+1}) - C(w^*) \leq \quad (22)$$

$$[\mathbb{E}C(w_t) - C(w^*)](1 - a_t m D) + AKM^2 a_t^2.$$

We are now in a position to state our result:

Theorem 3.4 *Let $C : \mathbb{R}^n \rightarrow \mathbb{R}$ be a twice differentiable cost function with unique minimum w^* . Assume that*

1. *Conditions C.1–C.6 from Theorem 3.2 hold with $B = 0$ in C.4.*
2. *Equation (20) holds.*
3. *$a_t = \frac{\tau}{t}$ with $\tau > 0$.*

Then $\mathbb{E}C(w_t) - \inf_{\tilde{w}} C(\tilde{w})$ is equivalent to $\frac{1}{t}$ as $t \rightarrow \infty$.

Proof Bottou and LeCun (2005, A.4) state: if $u_t = [1 - \frac{\alpha}{t} + o(1/t)]u_{t-1} + \frac{\beta}{t^2} + o(1/t^2)$ with $\alpha > 1$ and $\beta > 0$, then $tu_t \rightarrow \frac{\beta}{\alpha-1}$. Theorem 3.4 follows from (22) by setting $u_t = \mathbb{E}C(w_t) - C(w^*)$, noting that if (20) and hence (22) hold for one $D > 0$, they also hold for all larger constants. ■

4 Quasi-Newton Methods

Quasi-Newton methods are optimization methods with updates of the form $w_{t+1} \leftarrow w_t - a_t B_t \nabla_w C(w_t)$, where $a_t > 0$ is a scalar gain (typically set by a line search) and B_t a positive-semidefinite scaling matrix. If $B_t = I$, we have simple gradient descent; setting B_t to the inverse Hessian of $C(w)$ we recover Newton’s method. Inverting the $k \times k$ Hessian is a computationally challenging task if k is large; quasi-Newton methods reduce the computational cost by incrementally maintaining a symmetric positive-definite estimate B_t of the inverse Hessian of the objective function.

4.1 (L)BFGS

The BFGS algorithm (Nocedal and Wright, 1999) was developed independently by Broyden, Fletcher, Goldfarb, and Shanno in 1970. It incrementally maintains its estimate B_t of the inverse Hessian via a rank-two update that minimizes a weighted Frobenius norm $\|B_{t+1} - B_t\|_W$ subject to the *secant equation* $s_t = B_{t+1}y_t$, where $s_t := w_{t+1} - w_t$ and $y_t := \nabla_w C(w_{t+1}) - \nabla_w C(w_t)$ denote the most recent step along the optimization trajectory in parameter and gradient space, respectively. LBFGS is a limited-memory (matrix-free) version of BFGS.

4.2 Online (L)BFGS

Recently developed online variants of BFGS and LBFGS, called oBFGS *resp.* oLBFGS, are amenable to stochastic approximation of the gradient (Schraudolph et al., 2007). The key differences between the online and batch algorithms can be summarized as follows:

The gradient of the objective is estimated from small samples (mini-batches) of data. The difference y_t of gradients is computed with gradients for the same data sample, *i.e.*, for the same function $C(\cdot, z_t)$. Line search is replaced by a gain sequence a_t . A *trust region* parameter λ is introduced modifying the algorithm to estimate $(H_t + \lambda I)^{-1}$, where H_t is the Hessian at iteration t ; this prevents the largest eigenvalue of B_t from exceeding λ^{-1} .

See Schraudolph et al. (2007) for a more detailed description of the oLBFGS and oBFGS algorithms, and for experimental results on quadratic bowl objectives and conditional random fields (CRFs), which are instances of risk minimization.

Remark 4.1 *To guarantee a uniform lower eigenvalue bound for the updates of Schraudolph et al. (2007) we would have to use $B_t + \gamma I$ for some $\gamma > 0$, effectively interpolating between o(L)BFGS as defined by Schraudolph et al. (2007) and simple online gradient descent. This lower bound is not needed for convergence per se but to prove that the convergence is to the minimum.*

4.3 Filtering

Kailath et al. (2000, Chapter 14) present assumptions in control theory that imply either convergence of the matrices B_t or upper and lower eigenvalue bounds. The relevant control theory is too extensive and complicated to be reviewed here and we therefore only point out the connection.

5 Expert Advice with Bregman Divergences

We now compare the updates in the expert advice framework by Azoury and Warmuth (2001) to the SGD updates that would result from a non-linear reparametrization.

As outlined in section 2.3 the difference between the update

$$w_{t+1} = h^{-1}(h(w_t) - a_t \nabla L_t(w_t)). \quad (23)$$

and performing stochastic gradient descent with respect to a new variable $\gamma = h(w)$ is that the latter

would include matrices B_t which are the inverse Jacobians of h at w_t , i.e. the inverse Hessian of H where $\nabla H = h$. The update $\gamma_{t+1} = \gamma_t - a_t \nabla_{\gamma} \tilde{L}_t(\gamma_t)$ where $\tilde{L}_t(\gamma) = L_t(h^{-1}(\gamma))$, expressed with respect to w looks as follows

$$w_{t+1} = h^{-1}(h(w_t) - a_t B_t \nabla L_t(w_t)). \quad (24)$$

Therefore (23) can be written as

$$\gamma_{t+1} = \gamma_t - a_t B_t^{-1} \nabla_{\gamma} \tilde{L}_t(\gamma_t). \quad (25)$$

The main point of the reparametrization would be to change variables so that SGD converges with an optimal rate for the new objectives. We would like the new objective to be approximately quadratic. Assuming that we have chosen the transfer function in such a manner that our main theorems apply to the new objective function, it remains to check the scaling matrix condition.

To satisfy the conditions for the scaling matrices we need the Jacobian of the transfer function to have upper and lower eigenvalue bounds. In the literature that these methods are studied in, an assumption of uniformly bounded gradients $\nabla L_t(w_t)$ (Azoury and Warmuth, 2001) is often used, or the parameters are restricted to a compact set (Zinkevich, 2003). In that case the conditions on scaling matrices become easier to satisfy: in any compact set, popular functions like e^{θ} , $(1 + e^{\theta})^{-1}$, or other sigmoid functions (though not, e.g., θ^3 which is flat at the origin) have derivatives that are bounded from above and below by a strictly positive constant.

6 Conclusion

We provide a variable metric stochastic approximation theory which implies convergence for stochastic gradient descent even when the gradients are calculated with respect to variable metrics. Metrics are sometimes changed for optimization purposes, as in the case of using quasi-Newton methods. Our main theorems imply convergence results for online versions of the BFGS and LBFGS optimization methods. Kalman filters are a class of well-known algorithms that can be viewed as an online Newton method for the special case of square losses, since the procedure at every step performs a gradient step where the gradient is defined using the Hessian of the loss for the examples seen so far. Finally we investigate the task of learning from expert advice where Bregman divergences are frequently used to achieve updates that are suitable for the task at hand. We interpret the resulting updates as stochastic gradient descent in a space that has undergone a non-linear reparametrization, and where we use different metrics depending on the point we are at.

Acknowledgements

The authors are very grateful to Leon Bottou at NEC Labs in Princeton, NJ for his help with the main theorem.

The first author is funded by NICTA which is funded through the Australian Government's *Backing Australia's Ability initiative*, in part through the Australian Research Council.

References

- S. Amari and H. Nagaoka. *Methods of Information Geometry*. Oxford University Press, 1993.
- N. N. Schraudolph, J. Yu, and S. Günter. A stochastic quasi-Newton method for online convex optimization. In *Proc. 11th Intl. Conf. Artificial Intelligence and Statistics (AISTATS)*, San Juan, Puerto Rico, 2007. Society for Artificial Intelligence and Statistics. <http://nic.schraudolph.org/pubs/SchYuGue07.pdf>.
- L. Bottou and Y. LeCun. On-line learning for very large datasets. *Applied Stochastic Models in Business and Industry*, 21(2):137–151, 2005.
- S. Amari. Natural gradient works efficiently in learning. *Neural Computation*, 10(2):251–276, 1998.
- N. Murata. A statistical study of on-line learning. *Online learning in neural networks*, pages 63–92, 1998.
- J. Kivinen and M. K. Warmuth. Exponentiated gradient versus gradient descent for linear predictors. *Information and Computation*, 132(1):1–64, 1997.
- N. N. Schraudolph. Fast curvature matrix-vector products for second-order gradient descent. *Neural Computation*, 14(7):1723–1738, 2002.
- K. Azoury and M. K. Warmuth. Relative loss bounds for on-line density estimation with the exponential family of distributions. *Machine Learning*, 43(3):211–246, 2001. Special issue on Theoretical Advances in On-line Learning, Game Theory and Boosting.
- S. Shalev-Shwartz, Y. Singer, and N. Srebro. Pegasos: Primal estimated sub-gradient solver for SVM. In *Proc. Intl. Conf. Machine Learning*, 2007.
- E. Hazan, A. Agarwal, and S. Kale. Logarithmic regret algorithms for online convex optimization. *Machine Learning*, 69(2-3):169–192, 2007.
- M. Zinkevich. Online convex programming and generalised infinitesimal gradient ascent. In *Proc. Intl. Conf. Machine Learning*, pages 928–936, 2003.
- H. E. Robbins and S. Monro. A stochastic approximation method. *Annals of Mathematical Statistics*, 22: 400–407, 1951.

- J. R. Blum. Multidimensional stochastic approximation methods. *Annals of Mathematical Statistics*, 25: 737–744, 1954.
- H. E. Robbins and D. O. Siegmund. A convergence theorem for non negative almost supermartingales and some applications. In *Proc. Sympos. Optimizing Methods in Statistics*, pages 233–257, Ohio State Univ., Columbus, Ohio, 1971. Academic Press, New York.
- L. Bottou. Online algorithms and stochastic approximations. In D. Saad, editor, *Online Learning and Neural Networks*. Cambridge University Press, Cambridge, UK, 1998.
- J. Nocedal and S. J. Wright. *Numerical Optimization*. Springer Series in Operations Research. Springer, 1999.
- T. Kailath, A. Sayed, and B. Hassibi. *Linear Estimation*. Prentice Hall, 2000.